

#### Baseline results for Dataset FR1

	$D_\lambda^k$	$D_s^*$	$Q^*$
<b>EXP</b>	0.0174	0.2889	0.6987
<b>GS [1]</b>	0.2648	0.0113	0.7269
<b>GSA [2]</b>	0.0512	0.0121	0.9373
<b>AWLP [3]</b>	0.0274	0.0490	0.9249
<b>MTF-GLP [4]</b>	0.0396	0.0341	0.9277
<b>MF [5]</b>	0.0939	0.0663	0.8460

#### Baseline results for Dataset FR2

	$D_\lambda^k$	$D_s^*$	$Q^*$
<b>EXP</b>	0.0159	0.2415	0.7464
<b>GS [1]</b>	0.2073	0.0344	0.7654
<b>GSA [2]</b>	0.0555	0.0033	0.9414
<b>AWLP [3]</b>	0.0257	0.0367	0.9385
<b>MTF-GLP [4]</b>	0.0331	0.0254	0.9423
<b>MF [5]</b>	0.1044	0.0447	0.8556

#### Baseline results for Dataset RR1

	$Q2^n$	<i>SAM</i>	<i>ERGAS</i>
<b>EXP</b>	0.5740	3.3203	3.6816
<b>GS [1]</b>	0.5080	14.1040	7.9535
<b>GSA [2]</b>	0.6912	3.3297	2.8750
<b>AWLP [3]</b>	0.6166	4.8929	4.1890
<b>MTF-GLP [4]</b>	0.6207	3.6398	3.9223
<b>MF [5]</b>	0.6189	5.5063	7.2229

#### Baseline results for Dataset RR2

	$Q2^n$	<i>SAM</i>	<i>ERGAS</i>
<b>EXP</b>	0.5717	6.1252	7.0178
<b>GS [1]</b>	0.7221	5.5997	5.9123
<b>GSA [2]</b>	0.8251	4.8277	4.4092
<b>AWLP [3]</b>	0.6935	7.6157	5.9395
<b>MTF-GLP [4]</b>	0.7963	5.3705	4.7784
<b>MF [5]</b>	0.8057	5.2975	4.7005

The baseline results are obtained through the application of some state-of-the-art pansharpening methods belonging to the toolbox presented in [6]. The standard implementations can be downloaded by the following [link](#).

## References

- [1] C. A. Laben and B. V. Brower, "Process for enhancing the spatial resolution of multispectral imagery using pan-sharpening", Eastman Kodak Company, Tech. Rep. US Patent #6,011,875, 2000.
- [2] B. Aiazzi, S. Baronti and M. Selva, "Improving Component Substitution Pansharpening Through Multivariate Regression of MS+Pan Data," IEEE Trans. Geosci. Remote Sens., vol. 45, no. 10, pp. 3230-3239, Oct. 2007, doi: 10.1109/TGRS.2007.901007.
- [3] G. Vivone, L. Alparone, J. Chanussot, M. Dalla Mura, A. Garzelli, G. A. Licciardi, R. Restaino and L. Wald, "A critical comparison among pansharpening algorithms," IEEE Trans. Geosci. Remote Sens., vol. 53, no. 5, pp. 2565-2586, May 2015, doi: 10.1109/TGRS.2014.2361734.
- [4] B. Aiazzi, L. Alparone, S. Baronti, A. Garzelli and M. Selva, "MTF-tailored multiscale fusion of highresolution MS and Pan imagery," Photogramm. Eng. Remote Sens., vol. 72, no. 5, pp. 591–596, May 2006.
- [5] R. Restaino, G. Vivone, M. Dalla Mura and J. Chanussot, "Fusion of Multispectral and Panchromatic Images Based on Morphological Operators," IEEE Trans. Image Process., vol. 25, no. 6, pp. 2882-2895, June 2016, doi: 10.1109/TIP.2016.2556944.
- [6] G. Vivone, M. Dalla Mura, A. Garzelli, R. Restaino, G. Scarpa, M. O. Ulfarsson, L. Alparone and J. Chanussot, "A new benchmark based on recent advances in multispectral pansharpening: Revisiting pansharpening with classical and emerging pansharpening methods," IEEE Geosci. Remote Sens. Mag., vol. 9, no. 1, pp. 53-81, March 2021, doi: 10.1109/MGRS.2020.3019315.